Contents Volume 71, 1989

Preface
H. Okada (Fukuoka, Japan) and J.-P. Cadet (Paris, France) 1
Anatomy of trench-slope basins: examples from the Nankai Trough
H. Okada (Fukuoka, Japan) 3
Seaward jump of the subduction zone associated with ridge collision — An example from the Suruga Trough
K. Shimamura (Kumamoto, Japan) 15
Detailed topographic study of the Daiichi-Kashima Seamount
S. Tani (Tokyo, Japan) 31
Continental and marine influences expressed by deep-sea sedimentation off Japan (Kaiko Project)
H. Chamley and P. Debrabant (Villeneuve d'Ascq, France) 49
^{40}Ar — ^{39}Ar ages of igneous rocks recovered from Daiichi-Kashima and Erimo Seamounts during the Kaiko Project
Y. Takigani, I. Kaneoka, T. Ishii and Y. Nakamura (Tokyo, Japan) 71
Hydrochemistry in the Barbados Accretionary Complex: Leg 110 ODP
J. Gieskes (La Jolla, CA, U.S.A), G. Blanc (Paris, France), P. Vrolijk (Santa Cruz, CA, U.S.A.), J. C. Moore (Santa Cruz, CA, U.S.A.), A. Mascle (Rueil Malmaison, France), E. Taylor (College Station, TX, U.S.A.), P. Andrieff (Orléans, France), F. Alvarez (Palisades, NY, U.S.A.), R. Barnes (Anacortes, WA, U.S.A.), C. Beck (Villeneuve d'Ascq, France), J. Behrmann (Giessen, F.R.G.) K. Brown (Durham, England), M. Clark (Mobile, AL, U.S.A.), J. Dolan (Santa Cruz, CA, U.S.A.), A. Fisher (Miami, FL, U.S.A.), M. Hounslow (Sheffield, England), P. McLellan (Calgary, Canada), K. Moran (Dartmouth, Canada), Y. Ogawa (Fukuoka, Japan), T. Sakai (Utsunomiya, Japan), J. Schoonmaker (Honolulu, HI, U.S.A.), R. Wilkins (Cambridge, MA, U.S.A.) and C. Williams (Palisades, NY, U.S.A.) 83
Subduction-induced pore fluid venting and the formation of authigenic carbonates along the Cascadia continental margin: implications for the global Ca-cycle
M. W. Han and E. Suess (Corvallis, OR, U.S.A.) 97
Methane-derived CO_2 in pore fluids expelled from the Oregon subduction zone
E. Suess (Corvallis, OR, U.S.A.) and M. J. Whiticar (Hannover, F.R.G.) 119
Hinge structures, their variations and changes during growth, of some Japanese deep-sea, giant white clams, *Calyptogena*, collected during the "Kaiko" Project
M. Horikoshi (Chiba, Japan) 137
Adaptative features of the bivalve molluscs associated with fluid venting in the subduction zones off Japan
A. Fiala-Medioni (Banyuls-sur-Mer, France) and M. Le Pennec (Brest, France) 161
$^{13}C/^{12}C$ and $^{15}N/^{14}N$ ratios of vesicomyid clams and a vestimentiferan tube worm in the subduction zone east of Japan
T. Saino and S. Ohta (Tokyo, Japan) 169
Deep-sea communities dominated by the giant clam, *Calyptogena soyoae*, along the slope foot of Hatsushima Island, Sagami Bay, central Japan
J. Hashimoto (Yokosuka, Japan), S. Ohta (Tokyo, Japan), T. Tanaka (Yokosuka, Japan), H. Hotta (Yokosuka, Japan), S. Matsuzawa (Yokosuka, Japan) and H. Sakai (Tokyo, Japan) 179
Abyssal molluscan colony of *Calyptogena* in the Pliocene strata of the Miura Peninsula, central Japan
N. Niitsuma (Shizuoka, Japan), Y. Matsushima (Yokohama, Japan) and D. Hirata (Yokohama, Japan) 193
Euthecosomatous pteropods as paleohydrological and paleoecological indicators in a Tyrrhenian deep-sea core
J.W. Biekart (Utrecht, The Netherlands) 205
Paleoclimate and polar wander
W.L. Donn (Palisades, NY, U.S.A.) 225
Depositional environment and regional stratigraphic significance of the Serle Conglomerate: a Late Proterozoic submarine fan complex, South Australia
G.M. Young (London, Ont., Canada) and V.A. Gostin (Adelaide, S.A., Australia) 237
Distributional patterns along the Mediterranean continental margin (upper bathyal) using *Gryphus vitreus* (Brachiopoda) densities
C.C. Emig (Marseille, France) 253
Postglacial transgressions in Late Paleozoic basins of western Argentina: a record of glacioeustatic sea level rise
O.R. Lopez Gamundi (Buenos Aires, Argentina) 257
Relationships between climatically influenced sedimentation and salt diapirism in the French Western Alps based on evidence from organic and inorganic geochemistry
N.-P. Tribovillard and P. Cotillon (Villeurbanne, France) 271
Late Cretaceous pelagic sediments, volcanic ash and biotas from near the Louisville hotspot, Pacific Plate, paleolatitude ~42°S
P.F. Ballance (Auckland, New Zealand), J.A. Barron (Menlo Park, CA, U.S.A.), C.D. Blome (Denver, CO, U.S.A.), D. Bukry (La Jolla, CA, U.S.A.), P.A. Cawood (St. John's, Nfld, Canada), G.C.H. Chaproniere (Canberra, A.C.T., Australia), R. Frisch (Menlo Park, CA, U.S.A.), R.H. Herzer (Lower Hutt, New Zealand), C.S. Nelson (Hamilton,

New Zealand), P. Quinterno (Menlo Park, CA, U.S.A.), H. Ryan (Menlo Park, CA, U.S.A.), D.W. Scholl (Menlo Park, CA, U.S.A.), A.J. Stevenson (Menlo Park, CA, U.S.A.), D.G. Tappin (Nuku'alofa, Tonga) and T.L. Vallier (Menlo Park, CA, U.S.A.) 281

Tectonically-induced diversion of the Indus River west of the Salt Range, Pakistan
J.W. McDougall (Corvallis, OR, U.S.A.) 301

Predatory and non-predatory borings in echinoids from the upper Ocala Formation (Eocene), north-central Florida, U.S.A.
M.A. Gibson (Martin, TN, U.S.A.) and J.B. Watson (Knoxville, TN, U.S.A.). 309